

Construction of Supersymmetric Nonlinear Sigma Models on Noncompact Calabi-Yau Manifolds with Isometry

Kiyoshi Higashijima^{a*}, Tetsuji Kimura^a and Muneto Nitta^{b†}

^aDepartment of Physics, Graduate School of Science, Osaka University, Toyonaka, Osaka 560-0043, Japan

^bDepartment of Physics, Purdue University, West Lafayette, IN 47907-1396, USA

We propose a class of $\mathcal{N} = 2$ supersymmetric nonlinear sigma models on the noncompact Ricci-flat Kähler manifolds, interpreted as the complex line bundles over the hermitian symmetric spaces. Kähler potentials and Ricci-flat metrics for these manifolds with isometries are explicitly constructed by using the techniques of supersymmetric gauge theories. Each of the metrics contains a resolution parameter which controls the size of these base manifolds, and the conical singularity appears when the parameter vanishes.

1. Introduction

Nonlinear sigma models (NL σ Ms) in two dimensions are interesting for several reasons. They help us to understand various non-perturbative phenomena in four dimensional gauge theories such as confinement or dynamical mass generation. They also provide the description of strings propagating in the curved space-time. In the latter case, the consistency of strings requires the conformal invariance of the NL σ Ms. In the case of superstrings, furthermore, NL σ Ms have to possess the $\mathcal{N} = 2$ world sheet superconformal symmetry. Although all two-dimensional NL σ Ms with $\mathcal{N} = 2$ supersymmetry have $\mathcal{N} = 2$ superconformal symmetry at the tree level, they suffer from superconformal anomaly at the quantum level. Since these anomalies are proportional to the Ricci curvatures R_{ab} of the background (target) manifolds, $\mathcal{N} = 2$ supersymmetric NL σ Ms taking values on the target manifolds with vanishing Ricci curvatures, provide the possible candidates of the consistent superstrings propagating in the curved space-time. They are also candidates of finite quantum field theories in two

dimensions. $\mathcal{N} = 2$ supersymmetry in two dimensions requires the target manifolds of NL σ Ms are Kähler manifolds. The Kähler manifolds with vanishing Ricci curvatures are called the Calabi-Yau manifolds. Therefore, it is important to study the $\mathcal{N} = 2$ supersymmetric NL σ Ms defined on the Calabi-Yau manifolds.

To find the metric of the Ricci-flat manifolds, we have to solve the Einstein equation $R_{ab} = 0$ for the vacuum. If the manifold has enough symmetries, we can reduce the partial differential equation to an ordinary differential equation, which is usually easy to solve. In compact Calabi-Yau manifolds, however, there is no isometry and not a single explicit metric is known in this case. In some noncompact Calabi-Yau manifolds, the number of isometries is sufficient to reduce the Einstein equation to an ordinary differential equation. In this article, we report our recent progress to obtain the explicit metric for certain class of the noncompact Calabi-Yau manifolds [1,2,3].

2. NL σ Ms with $\mathcal{N} = 2$ Supersymmetry in Two Dimensions

$\mathcal{N} = 2$ supersymmetry requires that both scalars $A^a(x)$ and fermions $\psi^a(x)$ are complex fields. The scalar fields $A^a(x)$ ($a = 1, \dots, N$) are coordinates of the target manifold, whose

*Work supported in part by the Grant-in-Aid for Scientific Research

†Work supported by the U.S. Department of Energy under grant DE-FG02-91ER40681 (Task B)

metric is given by the Kähler potential $K(A, \bar{A})$ which characterizes the theory:

$$g_{a\bar{b}}(A, \bar{A}) = \frac{\partial^2 K(A, \bar{A})}{\partial A^a \partial \bar{A}^{\bar{b}}} = K_{,a\bar{b}}(A, \bar{A}). \quad (1)$$

The lagrangian for $\mathcal{N} = 2$ supersymmetric NL σ M is given by

$$\mathcal{L}(x) = g_{a\bar{b}} \partial_\mu A^a \partial^\mu \bar{A}^{\bar{b}} + i g_{a\bar{b}} \bar{\psi}^{\bar{b}} (\not{D} \psi)^a + \frac{1}{4} R_{a\bar{b}c\bar{d}} \psi^a \psi^c \bar{\psi}^{\bar{b}} \bar{\psi}^{\bar{d}}, \quad (2)$$

where the covariant derivative and the Riemann curvature are defined by

$$(D_\mu \psi)^a = \partial_\mu \psi^a + \partial_\mu A^b \Gamma_{bc}^a \psi^c \quad (3)$$

$$R_{a\bar{b}c\bar{d}} = \partial_c \partial_{\bar{d}} g_{a\bar{b}} - g^{e\bar{f}} \partial_c g_{a\bar{f}} \partial_{\bar{d}} g_{e\bar{b}}, \quad (4)$$

with the connection being

$$\Gamma_{ab}^c = g^{c\bar{d}} g_{b\bar{d},a} = g^{c\bar{d}} K_{,a\bar{d}\bar{b}}. \quad (5)$$

Since all geometrical quantities are derived from the Kähler potential K , all we have to do is to assume an appropriate symmetry of the scalar function K . Symmetries of a scalar function is far simpler than the symmetries of the tensor $g_{a\bar{b}}$.

3. Noncompact Calabi-Yau Manifolds

Although the NL σ M defined by (2) is conformally invariant at the tree level, it suffers from the conformal anomaly proportional to the Ricci curvature at the quantum level[4]. To obtain the conformally invariant theory at the quantum level, we impose the Ricci-flatness condition

$$R_{a\bar{b}} = 0 \quad (6)$$

to the target manifold.

In order to reduce an Einstein equation to an ordinary differential equation of single variable, we assume our target manifold M_1 has a symmetry, which carries each point of the manifold to a certain point on a line, which is parameterized by a single variable. In other word, any point of our manifold M_1 can be reached from a point on a line by a suitable action of some symmetry group G . If any point of the manifold can be reached from a single point, say the origin, by appropriate action of a symmetry group G , the manifold

is a homogeneous coset space G/H where H denotes a subgroup of G . The local structure of our manifold is, therefore, the product of a homogeneous space $M = G/H$ and a line. Since $\mathcal{N} = 2$ supersymmetry requires the manifold M_1 to be a Kähler manifold, we assume that the homogeneous space $M = G/H$ is *Kähler* by itself and the line is a *complex* line. Furthermore, we assume the Kähler coset space $M = G/H$ is a compact *Einstein manifold*, which satisfies the Einstein equation with a positive cosmological constant $h > 0$

$$R_{m\bar{n}} = h g_{m\bar{n}}, \quad (7)$$

where we use the indices m, n for the compact Kähler manifold $M = G/H$ whose complex dimension is $d = \dim(G/H)/2$. The complex dimension of the Ricci-flat manifold M_1 is $D = 1 + d$.

Our total space M_1 is parameterized by the complex coordinates ϕ_m ($m = 1, \dots, d$) and a complex number ρ that represents the complex line. We denote the Kähler potential of the compact Kähler-Einstein manifold M by $\Psi(\phi, \bar{\phi})$. Our assumption for the Kähler potential of the Ricci-flat manifold M_1 is as follows:

$$K(A, \bar{A}) = K(X), \quad (8)$$

where

$$X = \log |\rho|^2 + h \Psi(\phi, \bar{\phi}). \quad (9)$$

Now, if we notice

$$R_{a\bar{b}} = -g^{c\bar{d}} R_{a\bar{b}c\bar{d}} = -\partial_a \partial_{\bar{b}} \log \det(g_{c\bar{d}}) \quad (10)$$

the Ricci-flat condition (6) takes a very simple form

$$\det(g_{c\bar{d}}) = |\text{hol.}|^2, \quad (11)$$

where hol. is a holomorphic function of the coordinates ϕ and ρ .

Components of the metric for the whole space M_1

$$g_{a\bar{b}} = \begin{pmatrix} g_{\rho\bar{\rho}} & g_{\rho\bar{n}} \\ g_{m\bar{\rho}} & g_{m\bar{n}} \end{pmatrix}, \quad (12)$$

are given by

$$g_{\rho\bar{\rho}} = K'' \frac{\partial X}{\partial \rho} \frac{\partial X}{\partial \bar{\rho}}, \quad (13)$$

$$g_{\rho\bar{n}} = K'' \frac{\partial X}{\partial \rho} \frac{\partial X}{\partial \bar{\phi}^{\bar{n}}}, \quad (14)$$

$$g_{m\bar{n}} = K'' \frac{\partial X}{\partial \phi^m} \frac{\partial X}{\partial \bar{\phi}^{\bar{n}}} + K' \frac{\partial^2 X}{\partial \phi^m \partial \bar{\phi}^{\bar{n}}}, \quad (15)$$

where the prime denotes the differentiation with respect to the argument X . Using equations

$$\partial X / \partial \rho = 1/\rho (\rho \neq 0), \quad \frac{\partial^2 X}{\partial \phi^m \partial \bar{\phi}^{\bar{n}}} = h g_{m\bar{n}}, \quad (16)$$

we can express the determinant of the metric of M_1 in terms of the determinant of the metric of $M = G/H$

$$\det g_{a\bar{b}} = g_{\rho\bar{\rho}} \cdot \det(g_{m\bar{n}} - g_{\rho\bar{\rho}}^{-1} g_{m\bar{\rho}} g_{\rho\bar{n}}) \quad (17)$$

$$= \frac{1}{|\rho|^2} K'' (K')^d \cdot \det(h g_{m\bar{n}}). \quad (18)$$

Since we have assumed that the Kähler coset space is an Einstein space, we can easily calculate the determinant of its metric. By substituting the expressions of the Ricci curvature (10) and the metric $g_{m\bar{n}}$ in terms the Kähler potential Ψ (1) to the Einstein equation (7), we obtain

$$R_{m\bar{n}} = -\partial_m \partial_{\bar{n}} \log \det(g_{k\bar{l}}) = h \frac{\partial^2 \Psi(\phi, \bar{\phi})}{\partial \phi^m \partial \bar{\phi}^{\bar{n}}}, \quad (19)$$

which can be integrated to give a convenient formula for the determinant of the metric in Kähler-Einstein manifold M

$$\det(g_{m\bar{n}}) = e^{-h\Psi} |\text{hol.}|^2. \quad (20)$$

Combining the formula (18), (20) and (11), we obtain the Ricci-flatness condition for the Kähler potential of the manifold M_1

$$\det(g_{a\bar{b}}) = e^{-X} K'' (K')^d |\text{hol.}|^2 = (\text{const.}) \cdot |\text{hol.}|^2,$$

from which we find an ordinary differential equation for the Kähler potential:

$$e^{-X} \frac{d}{dX} (K')^D = a, \quad (21)$$

where a is a constant.

The equation (21) is easily solved to yield

$$K' = (\lambda e^X + b)^{\frac{1}{D}}, \quad (22)$$

where λ is a constant related to a and D , and b is an integration constant, interpreted as a

resolution parameter. To calculate the metric (13)(14)(15) of the Ricci-flat manifold M_1 , the Kähler potential itself is not necessary, though it is easily obtained by integrating (22).

Our method can be applied to any Kähler-Einstein coset manifold M , however, we need the Kähler potential Ψ of M to obtain the explicit expression for the metric of the noncompact Calabi-Yau manifold M_1 . Fortunately, the general prescription to calculate Ψ is given by Ito, Kugo and Kunitomo[5]. In this respect, our results gives the explicit realization of the method of Page and Pope, who constructed Ricci-flat manifolds by introducing a complex line as a fibre over a Kähler-Einstein manifold[6].

Here, we further require that the Kähler coset space M is a symmetric space, called the hermitian symmetric space, then the Kähler potential can be explicitly obtained by the gauge theory technique[2,7]. Therefore, we find a new class of Ricci-flat manifolds corresponding to various hermitian symmetric spaces with $h = C_2(G)/2$. For $M = CP^{N-1}$, M_1 reduces to the manifold obtained by Calabi[8].

REFERENCES

1. K. Higashijima, T. Kimura and M. Nitta, Phys. Lett. **B515** (2001) 421, [hep-th/0104184](#).
2. K. Higashijima, T. Kimura and M. Nitta, Phys. Lett. **B518** (2001) 301, [hep-th/0107100](#); Nucl. Phys. **B623** (2002) 133, [hep-th/0108084](#); Ann. Phys. **296** (2002) 347, [hep-th/0110216](#).
3. K. Higashijima, T. Kimura and M. Nitta, to appear in Nucl. Phys. B, [hep-th/0202064](#).
4. L. Alvarez-Gaumé and P. Ginsparg, Comm. Math. Phys. **102** (1985) 311.
5. K. Itoh, T. Kugo and H. Kunitomo, Nucl. Phys. **B263** (1986) 295; Prog. Theor. Phys. **75** (1986) 386.
6. D. N. Page and C. N. Pope, Class. Quantum Grav. **4** (1987) 213.
7. K. Higashijima and M. Nitta, Prog. Theor. Phys. **103** (2000) 635, [hep-th/9911139](#).
8. E. Calabi, Ann. Scient. Ec. Norm. Sup. **12** (1979) 269.